

6th Biennial Workshop on Japan-Kamchatka- Alaska Subduction Processes (JKASP-2009)



Mitigating natural hazards in active arc environments

Linkages among tectonism, earthquakes, magma genesis and eruption in volcanic arcs, with a special focus on hazards posed by arc volcanism and great earthquakes

JUNE 22-26, 2009

SCIENTIFIC PROGRAM &
ABSTRACTS

GEOPHYSICAL INSTITUTE
UNIVERSITY OF ALASKA FAIRBANKS
FAIRBANKS, ALASKA



UAF UNIVERSITY OF ALASKA
FAIRBANKS

MAJOR AND VOLATILE ELEMENTS IN GLASSES OF LARGE EXPLOSIVE ERUPTIONS IN KAMCHATKA CORRELATE WITH DEPTH TO SUBDUCTING PLATE

Kuvikas O.V.,^{1,4} Portnyagin M.V.,^{2,3} Ponomareva V.V.⁴

¹ *Kamchatka V. Bering State University, Petropavlovsk-Kamchatsky, Russia*

² *Institute of Geochemistry and Analytical Chemistry, Moscow, Russia*

³ *Leibniz Institute of Marine Sciences, IFM-GEOMAR, Kiel, Germany*

⁴ *Institute of Volcanology and Seismology, Petropavlovsk-Kamchatsky, Russia*

There were at least 40 large (>1 km³ of tephra) and numerous smaller explosive eruptions in Kamchatka during the Holocene. Tephra of strong eruptions have wide distribution and serve as time-markers in stratigraphic research. In previous studies, tephra of various volcanic centers were recognized based on K₂O content in bulk samples (Braitseva et al., 1997) and in volcanic glass (Ponomareva et al., 2004). In this study we attempted to assess multi-component systematics of volcanic glass from major Kamchatka tephra. These results are used for 1) fingerprinting of tephra from different sources; 2) comparison of silicic magmas across and along the volcanic arc, and 3) understanding the processes governing generation of silicic melts in Kamchatka.

Within the framework of the German-Russian project KALMAR, we have analyzed 94 samples of volcanic glass from major Kamchatka tephra. The analyses were obtained at the IFM-GEOMAR (Kiel, Germany) with the JEOL JXA 8200 microprobe using a single analytical protocol (15 keV accelerating voltage, 6 nA current, and 5 micron beam size). Smithsonian Institution glass standards (Jarosevich, 1980) were used for reference. High-sensitivity (H-type) crystals were used for analysis of volatiles and low concentration elements (Ca, Fe, Mg in silicic glasses), which significantly improved the analytical precision for these elements compared to conventional analyses. As a result we have obtained a new self-consistent database of ~2500 high-quality glass analyses of known source and age. Here we report some pilot results from the analysis of this database.

There is large compositional variability of volcanic glasses from different volcanic zones and volcanic centers in Kamchatka. Some spatial geochemical trends are, however, evident from our database. For example, mean glass compositions from different volcanoes demonstrate an increase in K₂O contents, and a decrease in FeO, MgO, CaO contents and Cl/K ratio from the volcanic front toward the rear-arc, i.e. with increasing depth to the subducting plate (Fig. 1). With regard to the increasing concentrations of incompatible elements (e.g., K) and decreasing Cl/K from the volcanic front to the rear-arc, the trends of silicic melts are qualitatively similar to those reported for mafic rocks and primitive melt inclusions in olivine (e.g., Volynets, 1994; Portnyagin et al., 2007) yet absolute concentrations of K₂O are higher in silicic melts and Cl/K ratios are lower. This suggests that crystal fractionation of mantle basaltic melts is the main mechanism of silicic melts generation in Kamchatka while assimilation of crustal rocks plays a relatively minor role. Reconciling the spatial geochemical zoning evident from our data and massive crustal assimilation involved in the origin of silicic melts (e.g., Bindeman et al., 2004) requires re-melting of rocks formed within the same volcanic center.

At first glance, most of volcanoes in Kamchatka produced tephra with glass compositions plotting along single fractionation trends. Clear exceptions from this pattern are Ksudach, Khangar, Shiveluch and Bezymianny volcanoes. Significant variability of glasses from individual volcanoes can be related to variable extent of crustal assimilation, compositional heterogeneity of parental melts and different conditions of crystallization. For example, glass from minor Bezymianny tephra erupted on October 14, 2007 differs dramatically from that found in the pumice from the 2400 yr BP large eruption by having lower Al₂O₃ and higher K₂O, FeO and TiO₂ at given SiO₂ content. We suggest that these compositional

peculiarities reflect low (near-surface) pressure conditions of magma fractionation in recent years (probably after 1956) that promoted frequent weak eruptions rather than the larger ones driven by high water pressure in deeper magma chamber.




Fig.1. Across-arc variations of major elements in volcanic glass. Symbols are labeled as follows: “SR” – Sredinny Range, “EVF” – Eastern Volcanic Front, “CKD” – Central Kamchatka Depression. Abbreviations for volcanoes: KO – Kurile Lake Caldera, AV – Avachinsky, KS – Ksudach, KRM – Karymskaya Caldera, ICH – Ichinsky, KHG – Khangar, KZ – Kizimen, OP – Opala, BZ – Bezymianny, SH – Shiveluch, TN-Taunshits, ZHP – Zhupanovskiy volcano, CHL – Chasha Lake, KK – Kekuknaiskiy volcano.

References

Braitseva, O.A., Ponomareva, V.V., Sulerzhitsky, L.D., Melekestsev, I.V., and Bailey, J. Holocene key-marker tephra layers in Kamchatka, Russia. *Quaternary Research*, 47, 125-139, 1997.

Jarosewich, E.J., Nelen, J.A., Norberg, J.A., Reference samples for electron microprobe analysis. *Geostandards Newsletter*, 4, 43-47, 1980.

Ponomareva, V., Bourgeois, J., Braitseva, O., Kozhurin, A., Kyle, P., Pinegina, T., Tibaldi, A., Major Holocene Marker Ash Layers on Kamchatka Peninsula, NW Pacific, and Their Implication for Dating Paleoseismic Events, *Eos Trans. AGU*, 85(47), Fall Meet. Suppl., Abstract V21, 07, 2004.

Portnyagin, M.V., Hoernle K., Plechov, P.Y., Mironov, N.L., Khubunaya, S.A., Constraints on mantle melting and composition and nature of slab components in volcanic arcs from volatiles (H₂O, S, Cl, F) and trace elements in melt inclusions from the Kamchatka Arc. *Earth and Planetary Science Letters* 255(1-2), 53-69, 2007.

Volynets, O.N., Geochemical types, petrology and genesis of Late Cenozoic volcanic rocks from the Kurile-Kamchatka island arc system. *Int. Geol. Rew.*, 36/4, 373-405, 1994.