

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/282656425>

Geology, petrology and geochemistry of the Tolbachik volcanic massif, Kamchatka, Russia

CONFERENCE PAPER · JUNE 2015

READS

12

14 AUTHORS, INCLUDING:



Boris Gordyechik

Russian Academy of Sciences

36 PUBLICATIONS 22 CITATIONS

[SEE PROFILE](#)



Osamu Ishizuka

National Institute of Advanced Industrial Sc...

138 PUBLICATIONS 1,434 CITATIONS

[SEE PROFILE](#)



T. Miyazaki

Japan Agency for Marine-Earth Science Tec...

65 PUBLICATIONS 493 CITATIONS

[SEE PROFILE](#)



Chiaki Toyama

Japan Agency for Marine-Earth Science Tec...

15 PUBLICATIONS 93 CITATIONS

[SEE PROFILE](#)



26th IUGG GENERAL ASSEMBLY 2015

INTERNATIONAL UNION OF GEODESY AND GEOPHYSICS
UNION GÉODÉSIQUE ET GÉOPHYSIQUE INTERNATIONALE

PRAGUE, CZECH REPUBLIC
PRAGUE CONGRESS CENTRE
JUNE 22–JULY 2, 2015

Earth and Environmental Sciences for Future Generations

VS28 Understanding VIPS (Volcanic and Igneous Plumbing Systems) through Multidisciplinary Research

IAVCEI (Volcanology,
Geochemistry)

VS28p-487

30-Jun-2015, 15:00 - 16:30



Abstract content:

Geology, petrology and geochemistry of the Tolbachik volcanic massif, Kamchatka, Russia

Data on the geology, petrography, and geochemistry of previously geochemically unstudied Middle-Late-Pleistocene rocks from Tolbachik volcanic massif (Central Kamchatka Depression, CKD) are presented. Two volcanic series – middle-K and high-K were erupted. The geochemical history of the massif was started earlier 86 ka (K-Ar dating) with the formation of the Tolbachik pedestal presented by middle-K series. During stratovolcanoes formation both series occur and the role of high-K melts was increasing with time. In Holocene high-K rocks are dominated but some cinder cone lavas are presented by middle-K high-Mg melts which suggest that both volcanic series are still exists. The computer modeling show that both series can be explained by the process of crystal fractionation at different water content from nearly or the same mantle source similar to high-Mg basalts of 1975 Northern Breakthrough. Middle-K rocks could crystallize at water-rich conditions (more than 2% of H₂O) while the high-K rock could crystallize at dry conditions at the same pressure. However the existence of different mantle sources and possible magma mixing cannot be excluded. Our data show that fractional crystallization at different P-T-H₂O-fO₂ conditions can be one of the main processes responsible for rock variations at CKD. Sr-Nd-Pb isotopes suggest 2-4% of crustal assimilation to the magma chamber during pedestal and stratovolcanoes formation while lava-cinder cones are not show evidences of crustal assimilation. Major and trace element data coupled with K-Ar dating provide strong evidence that Povorotnaya mount located in 8 km NE of Plosky Tolbachik is the old block of the Tolbachik massif pedestal and for the moment the oldest known object (306 ka by K-Ar dating) in Klyuchevskaya group.

Author(s):

T. Churikova¹, B. Gordychik², H. Iwamori³, H. Nakamura³, O. Ishizuka⁴, T. Nishizawa⁵, S. Haraguchi³, K. Yasukawa⁶, T. Miyazaki³, B. Vaglarov³, K. Ueki³, C. Toyama³, Q. Chang³, J.I. Kimura³.

¹Institute of Volcanology and Seismology, Far East Branch, Russian Academy of Sciences, Laboratory of Petrology and Geochemistry, Petropavlovsk-Kamchatsky, Russia.

²Institute of Experimental Mineralogy, Russian Academy of Sciences, Laboratory of Thermodynamics of Minerals, Chernogolovka, Russia.

³Japan Agency for Marine-Earth Science and Technology, Department of Solid Earth Geochemistry, Yokosuka, Japan.

⁴National Institute of Advanced Industrial Science and Technology, Institute of Geology and Geoinformation, Ibaraki, Japan.

⁵Tokyo Institute of Technology, Department of Earth and Planetary Sciences, Tokyo, Japan.

⁶University of Tokyo, Graduate School of Engineering, Tokyo, Japan.

Keywords:

Tolbachik Klyuchevskaya Group of Volcanoes subduction geochemistry fractional crystallization